



SpecialtyRx.GiantEagle.com  
1-844-259-1891

## Patient Information

New Patient  Current Patient

### Patient's Name

First \_\_\_\_\_ Last \_\_\_\_\_ MI \_\_\_\_\_

Male  Female

Last 4 digits of SSN \_\_\_\_\_ Date of Birth \_\_\_\_\_

Street Address \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ ZIP \_\_\_\_\_

Preferred Phone \_\_\_\_\_  Landline  Mobile

Alternate Phone \_\_\_\_\_  Landline  Mobile

Preferred Method of Contact  Call  Text

Email Address \_\_\_\_\_

Patient's Primary Language  English  Other If other, please specify \_\_\_\_\_

**Parent/Guardian Name (if under 18)** \_\_\_\_\_

Home Phone \_\_\_\_\_ Cell Phone \_\_\_\_\_

Email Address \_\_\_\_\_

**Alternate Caregiver/Contact** \_\_\_\_\_

OK to speak to/leave message with alternate caregiver/contact

Home Phone \_\_\_\_\_ Cell Phone \_\_\_\_\_

Email Address \_\_\_\_\_

**PLEASE ATTACH COPY OF FRONT AND BACK OF PATIENT'S INSURANCE CARD**



## Prescriber Information

Date Prescription Needed \_\_\_\_\_

Ship to Office  Patient Pickup at Retail  Ship to Home

Office Hours to Receive Shipment of Medication \_\_\_\_\_

Office Contact and Title \_\_\_\_\_

Office Contact Phone \_\_\_\_\_



## Prescribing Information Cont.

Medication	Strength	Directions	Qty/Refills
<input type="checkbox"/> Copaxone (glatiramer acetate)	<input type="checkbox"/> 20mg/mL prefilled syringe	Inject 20mg subcutaneously daily	Qty: <input type="checkbox"/> 30 syringes <input type="checkbox"/> 90 syringes Refills: _____
	<input type="checkbox"/> 40mg/mL prefilled syringe	Inject 40mg subcutaneously 3 times a week	Qty: <input type="checkbox"/> 12 syringes <input type="checkbox"/> 36 syringes Refills: _____
<input type="checkbox"/> Extavia (interferon beta-1b)	0.3mg vial	<input type="checkbox"/> Sig Titration Per Package Insert: <b>Weeks 1-2</b> inject 0.0625mg subcutaneously every other day <b>Weeks 3-4:</b> inject 0.125mg subcutaneously every other day <b>Weeks 5-6:</b> inject 0.1875mg subcutaneously every other day <b>Week 7 and thereafter:</b> inject 0.25mg subcutaneously every other day <input type="checkbox"/> Inject 0.25mg subcutaneously every other day	Qty: <input type="checkbox"/> 1 kit (15 single dose vials) <input type="checkbox"/> 3 kits (45 single dose vials) Refills: _____
<input type="checkbox"/> Gilenya (fingolimod)	0.5mg capsule	Take 1 capsule by mouth daily	Qty: <input type="checkbox"/> 30 capsules <input type="checkbox"/> 90 capsules Refills: _____
<input type="checkbox"/> Glatopa (glatiramer acetate)	<input type="checkbox"/> 20mg/mL prefilled syringe	Inject 20mg subcutaneously daily	Qty: <input type="checkbox"/> 30 syringes <input type="checkbox"/> 90 syringes Refills: _____
	<input type="checkbox"/> 40mg/mL prefilled syringe	Inject 40mg subcutaneously 3 times a week	Qty: <input type="checkbox"/> 12 syringes <input type="checkbox"/> 36 syringes Refills: _____
<input type="checkbox"/> Kesimpta (ofatumumab)	20mg/0.4mL auto-injector	<b>Starter:</b> <input type="checkbox"/> Inject 20mg subcutaneously once weekly for 3 doses at weeks 0, 1, and 2. Then start maintenance dose at week 4.	Qty: 3 auto-injectors Refills: 0
		<b>Maintenance:</b> <input type="checkbox"/> Inject 20mg subcutaneously once monthly	Qty: <input type="checkbox"/> 1 auto-injector <input type="checkbox"/> 3 auto-injectors Refills: _____

## Prescribing Information Cont.

Medication	Strength	Directions	Qty/Refills
<input type="checkbox"/> Plegridy (peginterferon beta-1a)	<b>Starter:</b> <input type="checkbox"/> 63mcg/0.5mL and 94mcg/0.5mL auto-injector kit <input type="checkbox"/> 63mcg/0.5mL and 94mcg/0.5mL prefilled syringe kit	<b>Starter:</b> Inject 63mcg subcutaneously on Day 1, inject 94mcg on day 15, then inject 125mcg on day 29 and every 14 days thereafter	Qty: 1 kit Refills: 0
	<b>Maintenance:</b> <input type="checkbox"/> 125mcg/0.5mL auto-injector <input type="checkbox"/> 125mcg/0.5mL subcutaneous prefilled syringe <input type="checkbox"/> 125mcg/0.5mL intramuscular prefilled syringe	<b>Maintenance:</b> <input type="checkbox"/> Inject 125mcg subcutaneously every 14 days <input type="checkbox"/> Inject 125mcg intramuscularly every 14 days	Qty: <input type="checkbox"/> 2 devices <input type="checkbox"/> 6 devices Refills: _____
<input type="checkbox"/> Rebif (interferon beta-1a)	<b>Starter:</b> <input type="checkbox"/> 6 x 8.8mcg and 6 x 22mcg prefilled syringes <b>starter kit</b> <input type="checkbox"/> 6 x 8.8mcg and 6 x 22mcg Rebidose auto-injector <b>starter kit (44mcg dose only)</b>	<b>Starter:</b> <input type="checkbox"/> Sig Titration for 22mcg dose ( <b>prefilled syringe only</b> ): <b>Weeks 1-2:</b> inject 4.4mcg subcutaneously 3 times per week <b>Weeks 3-4:</b> inject 11mcg subcutaneously 3 times per week <b>Weeks 5 and thereafter:</b> inject 22mcg subcutaneously 3 times per week <input type="checkbox"/> Sig Titration for 44mcg dose: <b>Weeks 1-2:</b> inject 8.8mcg subcutaneously 3 times per week <b>Weeks 3-4:</b> inject 22mcg subcutaneously 3 times per week <b>Weeks 5 and thereafter:</b> inject 44mcg subcutaneously 3 times per week	Qty: 1 starter kit Refills: 0
	<b>Maintenance:</b> <input type="checkbox"/> 22mcg Rebidose auto-injector <input type="checkbox"/> 44mcg Rebidose auto-injector <input type="checkbox"/> 22mcg prefilled syringe <input type="checkbox"/> 44mcg prefilled syringe	<input type="checkbox"/> Inject 22mcg subcutaneously 3 times per week <input type="checkbox"/> Inject 44mcg subcutaneously 3 times per week	Qty: <input type="checkbox"/> 12 devices <input type="checkbox"/> 36 devices Refills: _____
<input type="checkbox"/> Dimethyl Fumarate	<b>Starter:</b> <input type="checkbox"/> 120mg capsule	<b>Starter:</b> Take 120mg by mouth twice a day for 7 days	Qty: 14 capsules Refills: 0
	<b>Maintenance:</b> <input type="checkbox"/> 240mg capsule	<b>Maintenance:</b> Take 240mg by mouth twice a day	Qty: <input type="checkbox"/> 60 capsules <input type="checkbox"/> 180 capsules Refills: _____

## Prescribing Information Cont.

Medication	Strength	Directions	Qty/Refills
<input type="checkbox"/> Zeposia (ozanimod HCl)	<b>Starter:</b> <input type="checkbox"/> 4x 0.23mg capsules and 3x 0.46mg capsules ( <b>7 day starter kit</b> ) <input type="checkbox"/> 4x 0.23mg capsules, 3x 0.46mg capsules and 30x 0.92mg capsules ( <b>37 day starter kit</b> )	Take 0.23mg by mouth once daily on days 1 through 4, take 0.46mg on days 5 through 7, and then take 0.92mg once daily starting on day 8	Qty: <input type="checkbox"/> 1 starter kit (7 days) <input type="checkbox"/> 1 starter kit (37 days) Refills: 0
<input type="checkbox"/> Other: _____	<b>Maintenance:</b> <input type="checkbox"/> 0.92mg capsule	Take 0.92mg by mouth once daily	Qty: <input type="checkbox"/> 30 capsules <input type="checkbox"/> 90 capsules Refills: _____

Prescriber Name \_\_\_\_\_

Phone \_\_\_\_\_ Fax \_\_\_\_\_

Email Address \_\_\_\_\_

Office Address \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ ZIP \_\_\_\_\_

State License \_\_\_\_\_ DEA \_\_\_\_\_ NPI \_\_\_\_\_

In order for brand name to be dispensed, prescriber must hand write "Brand Medically Necessary" or "Brand Necessary" in the space below:

\_\_\_\_\_

I authorize this prescription and for Giant Eagle Specialty Pharmacy and its representatives to act as an agent to initiate and execute the insurance prior authorization process.

Prescriber signature required. NO STAMPS. Prescriber attests this is his/her legal signature.

Prescriber signature \_\_\_\_\_ Date \_\_\_\_\_